

Yang et al.

**PROCESS FOR MAKING PERFUME CONTAINING SURFACTANT
COMPOSITIONS HAVING PERFUME BURST AND ENHANCED
PERFUME DEPOSITION WHEN DILUTED**

FIELD OF THE INVENTION

The invention relates to a process for preparing compositions, e.g., personal wash perfume-containing compositions, which are able to deliver perfume benefits (i.e., “burst” or increase in perfume headspace relative to undiluted composition) upon dilution in water as well as enhanced deposition of one or more of the fragrance components on surfaces with which said compositions come into contact. More specifically, by being aware of which properties of the perfume and of the surfactant systems in which they are found are responsible for perfume release on dilution, applicants have been able to provide processes for selecting perfumes and/or surfactant systems in said compositions which meet defined criteria (as defined, for example, in perfume burst index) and thereby provide aroma burst of at least defined levels and enhanced perfume deposition upon dilution. The invention further relates to methods of changing fragrance note and/or introducing smell after dilution.

BACKGROUND OF THE INVENTION

Perfume is a key component for a favorable consumer experience with home and personal care products. It is also often the most costly component of the formulation. Typically, most of the fragrance is quickly lost as the product is used because most of the perfume is trapped in the surfactant system. There has been a long-standing need

to improve the utilization of perfumes and to design compositions that provide maximum and prolonged impact in use. It is one objective of this invention to develop a means by which products can be developed to provide greater fragrance impact and novel fragrance characteristics in use.

5

To this end, the applicant has found that perfume activity (e.g., the aroma of the perfume) both in a product (e.g., surfactant containing product) and upon use (e.g., dilution of product in shower) can be correlated to the thermodynamic characteristics of (1) the perfume itself and (2) the formulation in which the perfume is found.

10
15
20
25
30
35
40
45
50
55
60
65
70
75
80
85
90
95
100
105
110
115
120
125
130
135
140
145
150
155
160
165
170
175
180
185
190
195
200
205
210
215
220
225
230
235
240
245
250
255
260
265
270
275
280
285
290
295
300
305
310
315
320
325
330
335
340
345
350
355
360
365
370
375
380
385
390
395
400
405
410
415
420
425
430
435
440
445
450
455
460
465
470
475
480
485
490
495
500
505
510
515
520
525
530
535
540
545
550
555
560
565
570
575
580
585
590
595
600
605
610
615
620
625
630
635
640
645
650
655
660
665
670
675
680
685
690
695
700
705
710
715
720
725
730
735
740
745
750
755
760
765
770
775
780
785
790
795
800
805
810
815
820
825
830
835
840
845
850
855
860
865
870
875
880
885
890
895
900
905
910
915
920
925
930
935
940
945
950
955
960
965
970
975
980
985
990
995

For example, the degree to which a perfume will partition into oil or water (measured by a so-called "partition coefficient" and a reflection of the hydrophobicity of the perfume) and the degree to which the perfume evaporates (measured by "volatility constant" and a reflection of the volatility of the perfume) are two significant characteristics of the perfume which strongly affect the potential perfume "burst" when said product is diluted. By burst is meant an increase in the concentration of perfume in the vapor phase above the solution (i.e., this is often known as perfume headspace) with respect to the undiluted product and composition. The vapor phase can of course vary depending on product, for example, from a small area above a bottle of perfume to an area in a shower stall.

20

25

30

As noted, the perfume burst is affected not only by the perfume properties, but also by properties of the formulation in which the perfume is found. Thus, the number and type of surfactant micelles found in a surfactant solution also has an effect. For example, in a surfactant with high critical micellization concentration (CMC) (in compositions of high CMC, micelles do not form as readily) perfume "burst" would occur more readily and less dilution is required. The critical micellization concentration is defined as the surfactant concentration at which micelles begin to form from unassociated surfactant monomers (M.J. Rosen, Surfactants and Interfacial Phenomena, 2nd Ed., 1989). Conversely, in a surfactant with low critical micelle concentration (e.g., one where micelles do form easily or, stated differently, don't break

apart as readily once formed), a perfume, being generally more hydrophobic, tends to stay in the surfactant more readily. As a result, the perfume will tend not to "burst" (increase perfume headspace) as readily and, to achieve more headspace, more dilution may be required.

5

Other important formulation factors which may affect the "burst" of the perfume may include, but are not limited to, perfume content in solution and surfactant to water ratio.

Still another factor which can affect perfume "burst" is the environment in which it is found, e.g., such environmental factors as (1) overall sample amount; (2) vapor volume and (3) temperature.

According to the subject invention, applicants have succeeded in putting together a thermodynamic model which can be used to select the types of perfume and formulations which should be used in order to maximize this perfume burst or actual headspace (actual concentration of perfume in vapor phase) when a formulation (e.g., personal wash or shampoo formulation) is diluted in use. Also, applicants have found, since perfume burst correlates with perfume deposition, the model can be used to select perfumes in surfactant systems with enhanced deposition.

More specifically, applicants have defined a perfume burst index which defines compositions which can deliver a perfume burst (as well as enhanced deposition) upon dilution of at least a certain amount relative to undiluted composition; and further allows applicants to define a process for obtaining such compositions.

In general, the burst is achieved by diluting a surfactant system (e.g., an aqueous surfactant system) where burst begins upon dilution and maximum burst is obtained upon reaching CMC, therefore, releasing all of the perfume from the surfactant system. Thus, a composition yielding a maximum fragrance burst of 20% means the perfume concentration in the headspace increases by about 20% relative to the undiluted

product when the solution is diluted through the CMC. The CMC is the point where perfume-surfactant-water system changes to perfume-water system (i.e., system is too dilute for micelles to form).

5 A surfactant system is defined as a surfactant and/or surfactant mixtures which may include ingredients selected to manipulate the CMC in a continuous phase. These selected ingredients can include urea; glycerine; C₁-C₁₂ straight-chained or branched alcohols or diols; water soluble polymers such as polyvinylpyrrolidone, polyvinylalcohol, polyethyleneglycol, polypropyleneglycol; multivalent electrolytes such as magnesium, calcium and aluminum salts; and sugars such as dextrose, glucose, maltose, galactose, sucrose. The continuous phase is typically water, but may also include C₁-C₈ straight-chained or branched alcohols or diols, glycerine, C₁-C₈ esters and combinations thereof.

Generally, surfactants which may be used include, for example, anionic, nonionic, amphoteric/zwitterionic, and cationic surfactants as discussed in more detail below.

The subject invention relates to a process for preparing or selecting a composition having a fragrance burst, as measured by a "perfume burst index", of about 20% relative to a composition containing surfactant systems and perfume/fragrance prior to dilution of said product. The process is also to prepare or select a composition having enhanced deposition of one or more fragrance components on surface (e.g., skin) with which the composition comes into contact.

More specifically, the invention relates to a process for preparing specific compositions by selecting perfume and/or perfumes and surfactant systems and/or mixtures of surfactant systems and calculating therefrom a perfume burst index (PBI) according to the following formula:

$$\text{PBI} = \frac{\phi - 1.4/\text{CMC}}{K}$$

wherein ϕ = the oil/water partition coefficient of selected perfumes or perfumes components in a mixture;
 CMC = critical micellization concentration of surfactant systems or mixture of surfactant systems (wt./wt.);
 K = volatility constant of perfume or perfume components in a mixture from the continuous phase (atmospheres); and

The perfumes and surfactant systems are specifically selected to ensure that the PBI calculated is greater than about 3.

The process also defines a process to select fragrance or fragrances having enhanced deposition on surfaces onto which the fragrance(s) come into contact.

It should be understood that the PBI defines the maximum potential fragrance burst which is achieved at the CMC for the surfactant or surfactant mixture. For example, a relatively low PBI (e.g., about 3) will obtain a "burst" of at least 20% as noted. However, if the PBI is higher, much higher fragrance burst can be expected. Thus, for example, a burst of 20% may be achieved upon immediate dilution (assuming high enough PBI) and may continue to 700% or 800% or more at CMC (which as noted is point of maximum potential burst).

As far as applicants are aware, there is no art which specifically discloses that such burst can be obtained with such compositions or which discloses a way of predicting when and under what circumstances such fragrance "burst" will occur based on the dilution behavior of a perfume-surfactant-water system. Further no art of which applicants are aware discloses how such compositions are in turn related, for example, to properties of the perfume (e.g., partition coefficient, volatility) as well as to properties of the formulation (e.g., surfactant concentration and surfactant CMC).

BRIEF SUMMARY OF THE INVENTION

The invention relates to surfactant compositions, preferably aqueous surfactant compositions (e.g., bars, personal wash liquids, shampoos) in which perfume and surfactant are selected as to provide (1) a fragrance burst, when diluted, of at least about 20% relative to undiluted product; and (2) to have enhanced deposition of one or more fragrance components. Compositions must have a perfume burst index, as defined, of about 3.0 or greater, preferably 4.0 or greater.

The invention further relates to a process for selecting such compositions by selecting perfumes and choosing surfactant systems or manipulating CMC of surfactant system to ensure the PBI is at least about 3.0.

The invention is based on applicants' observation that variations in perfume impact ("burst") on dilution are essentially caused by the competition between a decrease in overall fragrance concentration upon dilution; and an increase in fragrance concentration in the continuous phase as the fragrance is released during disassociation of surfactant micelles which occurs during dilution. More specifically, and without wishing to be bound by theory, applicants have observed and shown that only perfumes with relatively large oil/water partition coefficient (very hydrophobic) possess the potential to produce a fragrance burst upon dilution. Other parameters which are

important to minimum burst include (1) volatility constant of perfume; (2) surfactant concentration; (3) type of surfactant system (e.g., CMC) etc.

Specifically, applicants have developed a theoretical mathematical equation
5 based on the various variables noted above. More specifically, they have developed a perfume burst index which both defines compositions having a maximum burst of at least 20% and further provides a process for selecting such compositions based on properties of perfume and surfactant system.

10 The relationships are defined in more detail below.

BRIEF DESCRIPTION OF THE INVENTION

Figure 1 is a chart showing various product parameters that influence perfume
5 performance in diluted products (e.g., diluted personal wash products).

Figure 2 shows theoretical calculation of fragrance burst for 3 types of perfume
(as per Table 1) with dilution.

Figure 3 shows fragrance burst profiles of different perfumes in surfactant
10 solution/

Figure 4 discloses an example of 2-component fragrances which would change
note upon dilution as one component increases in concentration and another
15 decreases.

Figure 5 shows theoretical models of fragrance burst with changes in surfactant
concentration.

Figure 6 shows experimental results of fragrance burst with changes in surfactant
20 concentration .

Figure 7 shows theoretical model of fragrance burst with change in surfactant
CMC.

Figure 8 shows experimental results of fragrance burst with change in surfactant
25 CMC.

Figure 9 shows normalized dilution curve for component in a perfume mixture.

Figure 10 shows result of sensory scores averaged over all panelists in a panel study of single perfume.

5 Figure 11 shows results of sensory scores averaged over all panelists in a panel study of multi-component perfume.

Figure 12 shows how deposition of perfume varies with CMC of surfactant.

10 Figure 13 shows deposition results for multi component perfume from surfactant systems having different CMCs.

DETAILED DESCRIPTION OF THE INVENTION

According to the subject invention, applicants have identified a process for preparing or selecting surfactant compositions having a defined fragrance burst by selecting perfume and/or perfumes and surfactant system and/or surfactant systems to provide said fragrance burst when diluted. The burst is at least 20% at dilution although it can be much higher and reaches maximum strength at CMC. Applicants have developed a process for determining (using defined perfume burst index) how to identify these compositions. That is, when a perfume is in a perfume-surfactant-water system (where some portion of perfume will be associated with surfactant micelles and some portion will be in water) and the formulation is diluted, a fragrance "burst" is produced when parameters are properly chosen. The intensity of the burst will depend, for example, on how much of the perfume is released from micelles as dilution occurs; and on the volatility of the perfume.

More specifically, some factors which affect perfume headspace concentration, upon dilution, are (a) various perfume properties (e.g., oil/water partition coefficient; ϕ , which is a measurement of the partitioning of the perfume or of the perfume components in a mixture between the oil and water phases thereby reflecting the hydrophobicity of perfume; and volatility constant, K , which is a measure of the partial pressure of the perfume or perfume components in a mixture in the vapor phase above the continuous phase of the composition, thereby reflecting the volatility of the perfume); (b) surfactant concentration in formulation (e.g., lower surfactant concentration tends to result in higher perfume headspace over initial formulation and more intense "burst" around CMC with fewer required dilutions); and (c) CMC of surfactant system (e.g., higher surfactant system CMC tends to provide more intense fragrance "burst" around CMC with fewer required dilutions).

As noted earlier, in general, for a fragrance "burst" or increase in perfume headspace concentration to occur during product use, the amount of fragrance released

from micelles as they dissociate must compensate for overall decrease in perfume concentration upon dilution. Compositions having a Perfume Burst Index (PBI) greater than about 3 clearly meet the necessary criteria.

5 Typically, a cleansing composition (e.g., a personal wash cleansing composition) containing perfume can be thought of as a perfume-oil-water system with the oil phase comprising surfactant micelles and water forming the continuous phase. Any perfume will distribute between the surfactant micelles and water phase as governed by the perfume's surfactant/water partition coefficient. In addition, the equilibrium perfume headspace concentration is proportional to the perfume concentration in the aqueous phase as governed by the volatility constant, K . The presence of a surfactant micelle phase impacts perfume headspace concentration by solubilizing the perfume and thereby affecting perfume concentration in the aqueous phase (e.g., more in micelles, less is in the water).

10 Use of, for example, a personal wash cleansing product in bath or shower causes dilution. This dilution leads to a decrease in micelle concentration forcing perfume into the aqueous phase and, accordingly, affecting the headspace concentration or perfume "burst".

15 20 However, dilution causes the quantity of surfactant micelles in a surfactant composition to decrease until gradually reaching critical micellization concentration (CMC). Below CMC, surfactant is present only as monomers or other small aggregates not large enough to form micelles. At this point, all perfume which was previously present in micelles is now totally released.

25 30 More specifically, it should be understood, CMC is the point of maximum potential burst (e.g., where release of perfume is complete because there are no more micelles). The release of perfume, however, begins as soon as dilution occurs and the burst gradually increases from first dilution up until CMC.

In measuring perfume headspace (using, for example, gas chromatography headspace measurement), applicants observed that dilution of certain compositions containing certain perfumes or certain perfume components in a perfume mixture led to an increase in perfume headspace and that, upon reaching the CMC, the maximum perfume headspace was achieved. It is precisely this "burst" or increase in headspace (e.g., from start of dilution up until CMC) and methods for determining in what compositions it will occur that is the basis of the subject invention.

The different parameters that affect perfume concentration in the vapor phase can be controlled by manipulating the formulation, the perfume itself or the environment in which the product dilution is occurring as shown in Figure 1. Understanding how each variable affects the overall fragrance performance in a personal product composition in use is critical to designing optimum systems for enhanced perfume benefits (e.g., fragrance burst, enhanced perfume deposition).

More specifically, applicants have found compositions which yield a maximum fragrance burst of at least about 20%, relative to undiluted product. These compositions are defined by establishing a perfume burst index (PBI) which is defined below:

$$\text{PBI} = \frac{\phi - 1.4/\text{CMC}}{K}$$

wherein ϕ = oil/water partition coefficient of perfume or perfume components in a mixture;
 CMC = critical micellization concentration (wt./wt.) of surfactant systems;
 K = volatility constant of perfume or perfume components in a mixture from the continuous phase (atmospheres);

wherein variables are selected such that PBI is greater than about 3.

As noted, perfume headspace behavior of a perfume-surfactant-water system is governed by different thermodynamic principles above or below CMC upon dilution. The perfume headspace behavior upon dilution is closely related to the oil/water partition coefficient (ϕ) and the volatility constant (K) of perfume, which reflect the hydrophobicity and volatility of the perfume respectively.

In Table 1 below, perfumes are classified into four categories according to their oil/water partition coefficients (ϕ) and their volatility constants (K). Since two of the three factors contributing to the PBI value for fragrance molecules are the partition coefficient and volatility constant one can predict the extent of "perfume burst" these different types of perfume should exhibit in the proper surfactant system.

Table 1: Classification of Perfume Molecules

	Examples of Partition Coefficient Φ (log Φ)	Hydrophobicity	Examples of Volatility Constant K^*	Volatility	Initial headspace	Burst upon dilution
For Perfume Type 1	50 (1.7)	Low	50	High	Very high	No
For Perfume Type 2	1000 (3)	High	1	Low	Very low	High
For Perfume Type 3	1000 (3)	High	250	High	High	Medium
For Perfume Type 4	50 (1.7)	Low	1	Low	Low	No

*K is generally measured in units of atmosphere.

Fragrance molecules that fall into the Type 1 category are actually very rare, but due to their low partition coefficient (ϕ) and their high volatility constant, one would predict no fragrance burst upon dilution. The low solubility in the surfactant phase would result in very small release of fragrance upon dilution, so a typical dilution profile (e.g., where perfume headspace decreases continuously with dilution) would be expected. That is, such molecules would normally tend to be found in the water phase anyway. So any action (dilution) causing release of perfume from surfactant micelles would be largely irrelevant.

In general, type 1 perfumes have oil/water partition coefficient (ϕ) of 1 to about 200, preferably 2 to 100; and volatility constant (K) of 20 to 1000, preferably 50 to 1000 atmospheres.

Type 2 perfumes would be anticipated to produce the most intense "perfume burst" upon dilution. These molecules are typically fragrance base notes and are very hydrophobic although their overall volatility is low resulting in low fragrance impact in a typical full formulation, i.e., before dilution. Therefore upon dilution, these molecules are released from the surfactant micelles and the noted increase in fragrance intensity is quite large. That is, because of their hydrophobicity, the perfume molecules tend to stay in micelles until release and that this release combined with the low volatility of the perfume in the initial formulation causes a high burst. In general, Type 2 perfumes have oil/water partition coefficients greater than 500, preferably greater than 700 and volatility constant less than about 20, preferably less than about 15.

Specific examples of Type 2 perfume molecules include allyl cyclohexane propionate, ambrettolide, Ambrox DL (dodecahydro-3a,6,6,9a-tetramethyl-naphtho[2,1-b]furan), amyl benzoate, amyl cinnamate, amyl cinnamic aldehyde, amyl salicylate, anethol, auranthol, benzophenone, benzyl butyrate, benzyl iso-valerate, benzyl salicylate, cadinene, campylcyclohexal, cedrol, cedryl acetate, cinnamyl cinnamate, citronellyl acetate, citronellyl isobutyrate, citronellyl propionate, cuminic aldehyde,

cyclohexylsalicylate, cyclamen aldehyde, cyclomyral, dihydro isojaconate, diphenyl methane, diphenyl oxide, dodecanal, dodecalactone, ethylene brassylate, ethylmethyl phenylglycidate, ethyl undecylenate, exaltolide, Galoxilide™ (1,3,4,6,7,8-hexahydro,4,6,6,7,8,8-hexamethyl-cyclopenta- γ -2-benzopyran), geranyl acetate, geranyl isobutyrate, hexadecanolide, hexenyl salicylate, hexyl cinnamic aldehyde, hexyl salicylate, α -ionone, β -ionone, γ -ionone, α -irone, isobutyl benzoate, isobutyl quinoline, Iso E Super™ (7-acetyl,1,2,3,4,5,6,7,8-octahydro,1,1,6,7-tetramethyl naphthalene), cis-jasmone, lillial, linalyl benzoate, 20 methoxy naphthalene, methyl cinnamate, methyl eugenol, γ -methylionone, methyl linolate, methyl linolenate, musk indanone, musk ketone, musk tibetine, myristicin, neryl acetate, δ -nonalactone, γ -nonalactone, patchouli alcohol, phantolide, phenylethyl benzoate, phenylethylphenylacetate, phenyl heptanol, phenyl hexanol, α -santalol, thibetolide, tonalid, δ -undecalactone, γ -undecalactone, vertenex, vetiveryl acetate, yara-yara and ylangene.

Hydrophobic, volatile fragrances fall into the Type 3 perfume category. Since these molecules have high oil/water partition coefficients, they will be solubilized by the surfactant system and then be released upon dilution providing a "perfume burst". The actual extent of this burst will not be as large as that for Type 2 perfumes because the fragrances are so volatile in the first place, hence the PBI values will be lower, but great enough to provide a perceivable burst. In general, they will have an oil/water partition coefficient of greater than about 500, preferably greater than 700; and volatility constant of about 20 to 1000.

Type 3 fragrance molecules fall into the top and middle note categories and include allo-ocimene, allyl caproate, allyl heptoate, anisole, camphene, carvacrol, carvone, citral, citronellal, citronellol, citronellyl nitrile, coumarin, cyclohexyl ethylacetate, p-cymene, decanal, dihydromyrcenol, dihydromyrcenyl acetate, dimethyl octanol, ethyllinalool, ethylhexyl ketone, eucalyptol, fenchyl acetate, geraniol, gernyl formate, hexenyl isobutyrate, hexyl acetate, hexyl neopentanoate, heptanal, isobornyl acetate, isoeugenol, isomenthone, isononyl acetate, isononyl alcohol, isomenthol, isopulegol,

limonene, linalool, linalyl acetate, menthyl acetate, methyl chavicol, methyl octyl
acetaldehyde, myrcene, naphthalene, nerol, neral, nonanal, 2-nonanone, nonyl acetate,
octanol, octanal, α -pinene, β -pinene, rose oxide, α -terpinene, γ -terpinene, α -terpinenol,
terpinolene, terpinyl acetate, tetrahydrolinalool, tetrahydromyrcenol, undecenal, veratrol,
and verdox.

Finally, Type 4 perfume components which again are classified as top and
middle notes, have very low oil/water partition coefficients and therefore like Type 1
components would not be expected to release significantly upon dilution to provide a
“perfume burst”. While these molecules will not add to the intensity of the fragrance
burst in a fully formulated perfume, they are still very important fragrance components.
By combining perfume molecules that decrease in intensity upon dilution with those that
“burst” upon dilution, the overall fragrance note during use will change. Fragrance note
is defined as the type of odor a perfume or mixture of perfume components produces.
For example, phenethylalcohol provides a floral rose note, methylionone produces a
woody note, and bergamot and limonene produces citrusy, fruity notes. The fragrance
could smell different in-use to that of the original product. Typically these perfumes will
have an oil/water partition coefficient of about 1 to about 200, preferably 2 to 100 and a
volatility constant of less than about 20, preferably less than about 15.

Fragrance Type 4 molecules include acetanisol; amyl acetate; anisic aldehyde;
anisylalcohol; benzaldehyde; benzyl acetate; benzyl acetone; benzyl alcohol; benzyl
formate; hexenol; laevo-carveol; d-carvone; cinnamaldehyde; cinnamic alcohol;
cinnamyl acetate; cinnamyl formate; cis-3-hexenyl acetate; Cyclal C (2,4-dimethyl-3-
cyclohexen-1-carbaldehyde); dihydroxyindole; dimethyl benzyl carbinol; ethyl acetate;
ethyl acetoacetate; ethyl butanoate; ethyl butyrate; ethyl vanillin; tricyclo decenyl
propionate; furfural; hexanal; hexenol; hydratropic alcohol; hydroxycitronellal; indole;
isoamyl alcohol; isopulegyl acetate; isoquinoline; ligustral; linalool oxide; methyl
acetophenone; methyl amyl ketone; methyl anthranilate; methyl benzoate; methyl
benzyl acetate; methyl heptenone; methyl heptyl ketone; methyl phenyl carbonyl acetate;
methyl salicylate; octalactone; para-cresol; para-methoxy acetophenone; para-methyl

acetophenone; phenethylalcohol; phenoxy ethanol; phenyl acetaldehyde; phenyl ethyl acetate; phenyl ethyl alcohol; prenyl acetate; propyl butyrate; safrole; vanillin and viridine.

5 In general, as noted above, it can be seen that molecules with higher oil/water partition coefficient (e.g., Type 2 and 3) will have a greater burst. This may be, without wishing to be bound by theory, because the perfume is highly soluble in the surfactant micelles and upon dilution, which triggers the dissociation of these micelles, a large amount of perfume is released thereby compensating for the decrease in overall perfume concentration in the diluted system. As noted from perfume Type 3 versus 2, 10 this is balanced to some extent by volatility (volatility constant) of the perfume since, for two perfumes otherwise having same oil/water partition coefficient, the one which is less volatile (Type 2) will be expected to have larger burst.

15 The CMC of the surfactant system also affects "perfume burst". Thus, for example, a surfactant system of relatively low CMC (e.g., sodium lauryl ether sulfate, also known as SLES) lowers the PBI and can greatly reduce the expected "perfume burst" on dilution. By contrast, surfactant system of relatively higher CMC (e.g., sodium laurate compared to SLES) would raise PBI and can thus increase "perfume burst" on 20 dilution. Examples of CMCs for various individual surfactants are set forth in Table 2 below.

Table 2: The Critical Micellization Concentration (CMC) of Surfactants

Surfactant	CMC (wt./wt.)
$C_8H_{17}SO_3^-Na^+$	0.0346*
$C_{10}H_{21}SO_3^-Na^+$	0.0105*
$C_{12}H_{25}SO_3^-Na^+$	0.0034*
$C_8H_{17}SO_4^-Na^+$	0.0325*
$C_{10}H_{21}SO_4^-Na^+$	0.00858*
$C_{11}H_{23}SO_4^-Na^+$	0.00438*
$C_{10}H_{21}OC_2SO_3^-Na^+$	0.00452*
$C_6H_{13}OOCCH_2SO_3^-Na^+$	0.0412*
$C_8H_{17}OOCCH_2SO_3^-Na^+$	0.0181*
$C_8H_{17}OOC(CH_2)_2SO_3^-Na^+$	0.0132*
$C_{12}H_{25}OOC(CH_2)_2SO_3^-Na^+$ (SCI)	0.00076* (30 °C)
$C_4H_9OOCCH_2CH(SO_3^-Na^+)COO C_4H_9$	0.0664*
$C_5H_{11}OOCH_2(SO_3^-Na^+)COO C_5H_{11}$	0.0178*
$C_6H_{13}OOCH_2CH(SO_3^-Na^+)COO C_6H_{13}$	0.0053*
$n-C_6H_{13}(OC_2H_4)_6OH$	0.0271*
$(C_2H_5)_2CHCH_2(OC_2H_4)_6OH$	0.0366*
$(C_3H_7)_2CHCH_2(OC_2H_4)_6OH$	0.0091*
$C_6H_{13}[OCH_2CH(CH_3)]_2(OC_2H_4)_{9.9}OH$	0.0307*
n-octyl- β -D-glucoside	0.0111*
Pluronic EO ₃₇ PO ₅₈ EO ₃₇	0.0065**
Sodium laureth sulfonate (2EO) (50°C)	0.0011***
Sodium laureth sulfonate (3EO) (50°C)	0.00084***
Linear alkylbenzene sulfonate	0.00171***
Sodium oleate	0.0008***
Potassium laurate	0.00607***
Sodium laurate	0.0052***
Caprylamidopropyl betaine	0.00032

Reference:

- 5 * M. J. Rosen, Surfactants and Interfacial Phenomena, 2nd ed., 1989
 ** P. Alexandridis, L. Yang, Macromolecules, 2000, 33, 5574-5587.
 *** P. Mukerjee, K. J. Mysels, Critical Micelle Concentrations of Aqueous Surfactant Systems, 1971

The CMC of the surfactant or surfactant system mixture can itself be manipulated (e.g., to help control “perfume burst”) by using various components which can be added to affect the CMC.

5

Non-limiting examples of such component include urea; glycerine; C_1 - C_{12} straight-chained or branched alcohols or diols; water soluble polymers such as polyvinylpyrrolidone, polyvinylalcohol polyethylene glycol, or polypropyleneglycol; multivalent electrolytes (e.g., magnesium, calcium and aluminum salts); and sugars such as dextrose, glucose, maltose, galactose and sucrose.

10
11
12
13
14
15
16
17
18
19
20
21
22
23
24
25
26
27
28
29
30
31
32
33
34
35
36
37
38
39
40
41
42
43
44
45
46
47
48
49
50
51
52
53
54
55
56
57
58
59
60
61
62
63
64
65
66
67
68
69
70
71
72
73
74
75
76
77
78
79
80
81
82
83
84
85
86
87
88
89
90
91
92
93
94
95
96
97
98
99
100
101
102
103
104
105
106
107
108
109
110
111
112
113
114
115
116
117
118
119
120
121
122
123
124
125
126
127
128
129
130
131
132
133
134
135
136
137
138
139
140
141
142
143
144
145
146
147
148
149
150
151
152
153
154
155
156
157
158
159
160
161
162
163
164
165
166
167
168
169
170
171
172
173
174
175
176
177
178
179
180
181
182
183
184
185
186
187
188
189
190
191
192
193
194
195
196
197
198
199
200
201
202
203
204
205
206
207
208
209
210
211
212
213
214
215
216
217
218
219
220
221
222
223
224
225
226
227
228
229
230
231
232
233
234
235
236
237
238
239
240
241
242
243
244
245
246
247
248
249
250
251
252
253
254
255
256
257
258
259
260
261
262
263
264
265
266
267
268
269
270
271
272
273
274
275
276
277
278
279
280
281
282
283
284
285
286
287
288
289
290
291
292
293
294
295
296
297
298
299
300
301
302
303
304
305
306
307
308
309
310
311
312
313
314
315
316
317
318
319
320
321
322
323
324
325
326
327
328
329
330
331
332
333
334
335
336
337
338
339
340
341
342
343
344
345
346
347
348
349
350
351
352
353
354
355
356
357
358
359
360
361
362
363
364
365
366
367
368
369
370
371
372
373
374
375
376
377
378
379
380
381
382
383
384
385
386
387
388
389
390
391
392
393
394
395
396
397
398
399
400
401
402
403
404
405
406
407
408
409
410
411
412
413
414
415
416
417
418
419
420
421
422
423
424
425
426
427
428
429
430
431
432
433
434
435
436
437
438
439
440
441
442
443
444
445
446
447
448
449
450
451
452
453
454
455
456
457
458
459
460
461
462
463
464
465
466
467
468
469
470
471
472
473
474
475
476
477
478
479
480
481
482
483
484
485
486
487
488
489
490
491
492
493
494
495
496
497
498
499
500
501
502
503
504
505
506
507
508
509
510
511
512
513
514
515
516
517
518
519
520
521
522
523
524
525
526
527
528
529
530
531
532
533
534
535
536
537
538
539
540
541
542
543
544
545
546
547
548
549
550
551
552
553
554
555
556
557
558
559
560
561
562
563
564
565
566
567
568
569
570
571
572
573
574
575
576
577
578
579
580
581
582
583
584
585
586
587
588
589
590
591
592
593
594
595
596
597
598
599
600
601
602
603
604
605
606
607
608
609
610
611
612
613
614
615
616
617
618
619
620
621
622
623
624
625
626
627
628
629
630
631
632
633
634
635
636
637
638
639
640
641
642
643
644
645
646
647
648
649
650
651
652
653
654
655
656
657
658
659
660
661
662
663
664
665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702
703
704
705
706
707
708
709
710
711
712
713
714
715
716
717
718
719
720
721
722
723
724
725
726
727
728
729
730
731
732
733
734
735
736
737
738
739
740
741
742
743
744
745
746
747
748
749
750
751
752
753
754
755
756
757
758
759
760
761
762
763
764
765
766
767
768
769
770
771
772
773
774
775
776
777
778
779
780
781
782
783
784
785
786
787
788
789
790
791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811
812
813
814
815
816
817
818
819
820
821
822
823
824
825
826
827
828
829
830
831
832
833
834
835
836
837
838
839
840
841
842
843
844
845
846
847
848
849
850
851
852
853
854
855
856
857
858
859
860
861
862
863
864
865
866
867
868
869
870
871
872
873
874
875
876
877
878
879
880
881
882
883
884
885
886
887
888
889
890
891
892
893
894
895
896
897
898
899
900
901
902
903
904
905
906
907
908
909
910
911
912
913
914
915
916
917
918
919
920
921
922
923
924
925
926
927
928
929
930
931
932
933
934
935
936
937
938
939
940
941
942
943
944
945
946
947
948
949
950
951
952
953
954
955
956
957
958
959
960
961
962
963
964
965
966
967
968
969
970
971
972
973
974
975
976
977
978
979
980
981
982
983
984
985
986
987
988
989
990
991
992
993
994
995
996
997
998
999
1000
1001
1002
1003
1004
1005
1006
1007
1008
1009
1010
1011
1012
1013
1014
1015
1016
1017
1018
1019
1020
1021
1022
1023
1024
1025
1026
1027
1028
1029
1030
1031
1032
1033
1034
1035
1036
1037
1038
1039
1040
1041
1042
1043
1044
1045
1046
1047
1048
1049
1050
1051
1052
1053
1054
1055
1056
1057
1058
1059
1060
1061
1062
1063
1064
1065
1066
1067
1068
1069
1070
1071
1072
1073
1074
1075
1076
1077
1078
1079
1080
1081
1082
1083
1084
1085
1086
1087
1088
1089
1090
1091
1092
1093
1094
1095
1096
1097
1098
1099
1100
1101
1102
1103
1104
1105
1106
1107
1108
1109
1110
1111
1112
1113
1114
1115
1116
1117
1118
1119
1120
1121
1122
1123
1124
1125
1126
1127
1128
1129
1130
1131
1132
1133
1134
1135
1136
1137
1138
1139
1140
1141
1142
1143
1144
1145
1146
1147
1148
1149
1150
1151
1152
1153
1154
1155
1156
1157
1158
1159
1160
1161
1162
1163
1164
1165
1166
1167
1168
1169
1170
1171
1172
1173
1174
1175
1176
1177
1178
1179
1180
1181
1182
1183
1184
1185
1186
1187
1188
1189
1190
1191
1192
1193
1194
1195
1196
1197
1198
1199
1200
1201
1202
1203
1204
1205
1206
1207
1208
1209
1210
1211
1212
1213
1214
1215
1216
1217
1218
1219
1220
1221
1222
1223
1224
1225
1226
1227
1228
1229
1230
1231
1232
1233
1234
1235
1236
1237
1238
1239
1240
1241
1242
1243
1244
1245
1246
1247
1248
1249
1250
1251
1252
1253
1254
1255
1256
1257
1258
1259
1260
1261
1262
1263
1264
1265
1266
1267
1268
1269
1270
1271
1272
1273
1274
1275
1276
1277
1278
1279
1280
1281
1282
1283
1284
1285
1286
1287
1288
1289
1290
1291
1292
1293
1294
1295
1296
1297
1298
1299
1300
1301
1302
1303
1304
1305
1306
1307
1308
1309
1310
1311
1312
1313
1314
1315
1316
1317
1318
1319
1320
1321
1322
1323
1324
1325
1326
1327
1328
1329
1330
1331
1332
1333
1334
1335
1336
1337
1338
1339
1340
1341
1342
1343
1344
1345
1346
1347
1348
1349
1350
1351
1352
1353
1354
1355
1356
1357
1358
1359
1360
1361
1362
1363
1364
1365
1366
1367
1368
1369
1370
1371
1372
1373
1374
1375
1376
1377
1378
1379
1380
1381
1382
1383
1384
1385
1386
1387
1388
1389
1390
1391
1392
1393
1394
1395
1396
1397
1398
1399
1400
1401
1402
1403
1404
1405
1406
1407
1408
1409
1410
1411
1412
1413
1414
1415
1416
1417
1418
1419
1420
1421
1422
1423
1424
1425
1426
1427
1428
1429
1430
1431
1432
1433
1434
1435
1436
1437
1438
1439
1440
1441
1442
1443
1444
1445
1446
1447
1448
1449
1450
1451
1452
1453
1454
1455
1456
1457
1458
1459
1460
1461
1462
1463
1464
1465
1466
1467
1468
1469
1470
1471
1472
1473
1474
1475
1476
1477
1478
1479
1480
1481
1482
1483
1484
1485
1486
1487
1488
1489
1490
1491
1492
1493
1494
1495
1496
1497
1498
1499
1500
1501
1502
1503
1504
1505
1506
1507
1508
1509
1510
1511
1512
1513
1514
1515
1516
1517
1518
1519
1520
1521
1522
1523
1524
1525
1526
1527
1528
1529
1530
1531
1532
1533
1534
1535
1536
1537
1538
1539
1540
1541
1542
1543
1544
1545
1546
1547
1548
1549
1550
1551
1552
1553
1554
1555
1556
1557
1558
1559
1560
1561
1562
1563
1564
1565
1566
1567
1568
1569
1570
1571
1572
1573
1574
1575
1576
1577
1578
1579
1580
1581
1582
1583
1584
1585
1586
1587
1588
1589
1590
1591
1592
1593
1594
1595
1596
1597
1598
1599
1600
1601
1602
1603
1604
1605
1606
1607
1608
1609
1610
1611
1612
1613
1614
1615
1616
1617
1618
1619
1620
1621
1622
1623
1624
1625
1626
1627
1628
1629
1630
1631
1632
1633
1634
1635
1636
1637
1638
1639
1640
1641
1642
1643
1644
1645
1646
1647
1648
1649
1650
1651
1652
1653
1654
1655
1656
1657
1658
1659
1660
1661
1662
1663
1664
1665
1666
1667
1668
1669
1670
1671
1672
1673
1674
1675
1676
1677
1678
1679
1680
1681
1682
1683
1684
1685
1686
1687
1688
1689
1690
1691
1692
1693
1694
1695
1696
1697
1698
1699
1700
1701
1702
1703
1704
1705
1706
1707
1708
1709
1710
1711
1712
1713
1714
1715
1716
1717
1718
1719
1720
1721
1722
1723
1724
1725
1726
1727
1728
1729
1730
1731
1732
1733
1734
1735
1736
1737
1738
1739
1740
1741
1742
1743
1744
1745
1746
1747
1748
1749
1750
1751
1752
1753
1754
1755
1756
1757
1758
1759
1760
1761
1762
1763
1764
1765
1766
1767
1768
1769
1770
1771
1772
1773
1774
1775
1776
1777
1778
1779
1780
1781
1782
1783
1784
1785
1786
1787
1788
1789
1790
1791
1792
1793
1794
1795
1796
1797
1798
1799
1800
1801
1802
1803
1804
1805
1806
1807
1808
1809
1810
1811
1812
1813
1814
1815
1816
1817
1818
1819
1820
1821
1822
1823
1824
1825
1826
1827
1828
1829
1830
1831
1832
1833
1834
1835
1836
1837
1838
1839
1840
1841
1842
1843
1844
1845
1846
1847
1848
1849
1850
1851
1852
1853
1854
1855
1856
1857
1858
1859
1860
1861
1862
1863
1864
1865
1866
1867
1868
1869
1870
1871
1872
1873
1874
1875
1876
1877
1878
1879
1880
1881
1882
1883
1884
1885
1886
1887
1888
1889
1890
1891
1892
1893
1894
1895
1896
1897
1898
1899
1900
1901
1902
1903
1904
1905
1906
1907
1908
1909
1910
1911
1912
1913
1914
1915
1916
1917
1918
1919
1920
1921
1922
1923
1924
1925
1926
1927
1928
1929
1930
1931
1932
1933
1934
1935
1936
1937
1938
1939
1940
1941
1942
1943
1944
1945
1946
1947
1948
1949
1950
1951
1952
1953
1954
1955
1956
1957
1958
1959
1960
1961
1962
1963
1964
1965
1966
1967
1968
1969
1970
1971
1972
1973
1974
1975
1976
1977
1978
1979
1980
1981
1982
1983
1984
1985
1986
1987
1988
1989
1990
1991
1992
1993
1994
1995
1996
1997
1998
1999
2000
2001
2002
2003
2004
2005
2006
2007
2008
2009
2010
2011
2012
2013
2014
2015
2016
2017
2018
2019
2020
2021
2022
2023
2024
2025
2026
2027
2028
2029
2030
2031
2032
2033
2034
2035
2036
2037
2038
2039
2040
2041
2042
2043
2044
2045
2046
2047
2048
2049
2050
2051
2052
2053
2054
2055
2056
2057
2058
2059
2060
2061
2062
2063
2064
2065
2066
2067
2068
2069
2070
2071
2072
2073
2074
2075
2076
2077
2078
2079
2080
2081
2082
2083
2084
2085
2086
2087
2088
2089
2090
2091
2092
2093
2094
2095
2096
2097
2098
2099
2100
2101
2102
2103
2104
2105
2106
2107
2108
2109
2110
2111
2112
2113
2114
2115
2116
2117
2118
2119
2120
2121
2122
2123
2124
2125
2126
2127
2128
2129
2130
2131
2132
2133
2134
2135
2136
2137
2138
2139
2140
2141
2142
2143
2144
2145
2146
2147
2148
2149
2150
2151
2152
2153
2154
2155
2156
2157
2158
2159
2160
2161
2162
2163
2164
2165
2166
2167
2168
2169
2170
2171
2172
2173
2174
2175
2176
2177
2178
2179
2180
2181
2182
2183
2184
2185
2186
2187
2188
2189
2190
2191
2192
2193
2194
2195
2196
2197
2198
2199
2200
220

Zwitterionic surfactants can be broadly described as derivatives of aliphatic quaternary ammonium, phosphonium and sulfonium compounds in which aliphatic radicals can be straight or branched chain and wherein at least one aliphatic substituent contains about 8 to about 18 carbons and at least one contains an anionic group, e.g., carboxy, sulfonate, sulfate, phosphate or phosphonate.

Amphoteric surfactants include at least one acid group (e.g., carboxylic or sulfonic). They include quaternary nitrogen and may include quaternary amido acids as acid group. They also generally include alkyl or alkenyl group of 7 to 18 carbons. Example include simple betaines, amido betaines and sulfobetaines.

Nonionic surfactants which may be used include reaction product of compounds having a hydrophobic group and a reactive hydrogen (for example, aliphatic alcohols, acids, amides or alkyl phenols) with alkylene oxide, especially ethylene oxide either alone or with propylene oxide. Examples include alkyl phenols-ethylene oxide condensates and condensation products of aliphatic (C₈-C₁₈) primary or secondary linear or branched alcohols with ethylene oxide. They may also be sugar amides and alkylpolyglycosides.

Cationic surfactants include quaternary ammonium compounds such as, for example, alkyldimethyl ammonium halogenides.

Some examples of surfactants which may be used in surfactant systems of the invention include sodium lauryl ether sulfate, sodium laurate, potassium lauride, sodium oleate, potassium oleate, triethanolamine laurate, alkylpolyglucosides, sodium lauryl sulfate, caprylamidopropyl betaine and sodium cocoylisethionate. As noted, however, there is theoretically no limit as to which surfactant or surfactant systems may be used.

The compositions of the invention also have enhanced deposition of perfume or perfume components onto the surfaces with which the compositions come into contact.

Generally, for compositions with higher PBI, perfume headspace increases with dilution due to higher perfume concentration in the continuous phase. The higher concentration in the continuous phase also favors greater surface contact and hence greater perfume deposition. Thus, perfume deposition is correlated to perfume burst.

5 In this way, it can be seen that PBI can be used to predict fragrance delivery or deposition. Those surfactant systems which facilitate perfume burst will, in other words, have enhanced deposition relative to surfactant systems which do not favor burst as much.

10 In another embodiment, the invention relates to a method of changing a fragrance note in a composition subsequent to dilution by (1) selecting a fragrance with a mixture of perfume components wherein some have PBI greater than 3.0 and some have PBI less than 3.0; and (2) assuring selection of fragrance components such that the components desired in the new fragrance (i.e., notes with desired fragrance after dilution) have a PBI greater than about 3.

15 In yet another embodiment, the invention relates to a method of introducing a desired fragrance to a composition subsequent to dilution by ensuring that fragrance or fragrance components having the desired aroma and which are selected for the composition, have a PBI greater than about 3 so that, upon dilution, the perfume headspace concentration exceeds the odor threshold. Odor threshold is the minimum concentration of fragrance in the vapor phase required for human detection.

20 Except in the operating and comparative examples, or where otherwise explicitly indicated, all numbers in this description indicating amounts or ratios of materials or conditions or reaction, physical properties of materials and/or use are to be understood as modified by the word "about".

Where used in the specification, the term “comprising” is intended to include the presence of stated features, integers, steps, components, but not to preclude the presence or addition of one or more features, integers, steps, components or groups thereof.

The following examples are intended to further illustrate the invention and are not intended to limit the invention in any way.

Unless indicated otherwise, all percentages are intended to be percentages by weight.

EXAMPLES

EXAMPLE 1

Fragrance molecules vary quite a bit in their physical properties and can therefore provide a variety of different effects depending on properties such as hydrophobicity, water solubility, volatility, molecular weight, etc. As shown in Example 1, some perfume components with specific physical parameters will provide a “fragrance burst” upon dilution of surfactant containing products such as shampoos, shower gels, soap bars, etc. A “perfume burst index” has been developed to determine which fragrance molecules will provide a burst or an increase in perfume in the air upon dilution of the product (which typically happens during standard use of shampoos, shower gels, soap bars, light duty liquids, etc.). This “perfume burst index” (PBI) is a function of several physical properties of the perfume and of the surfactant system. Depending on the amount of fragrance burst one desires, a certain PBI is required.

$$PBI = \frac{\phi - 1.4 / CMC}{K}$$

ϕ = oil/water partition coefficient of selected perfume or perfume component;

CMC = critical micellization concentration (wt/wt) of surfactant systems;

K = volatility constant of perfume or perfume components of a mixture from the continuous phase

To achieve a small, but potentially noticeable maximum fragrance burst of 20% from a product containing surfactant, the PBI of the perfume should be greater than about 3.0. To produce a 50% enhancement of the fragrance, the PBI needs to be greater than about 11 and to double the amount of fragrance upon use, the PBI should be greater than about 27. A “perfume burst index” can be calculated for any desired perfume molecule in a surfactant system. Table 3 contains a list of fragrance molecules and their PBI’s in a product with CMC of 0.005 wt/wt (i.e. sodium laurate).

Table 3: PBI Values for Various Perfumes

Perfume Molecule	PBI	"20% Burst"?
Acetanisol	-31,000	No
p-Cresol	-27,778	No
Indole	-5,209	No
Benzyl Acetate	-819	No
2-Phenoxyethanol	-156	No
2-heptanone	-153	No
Acetaldehyde	-130	No
Cinnamyl Acetate	-122	No
Hexanal	-109	No
Methyl Benzoate	-13	No
Anisole	-5	No
Limonene	12	Yes
p-Cymene	12.2	Yes
α -pinene	20	Yes
Isoeugenol	58	Yes
Menthone	61	Yes
Napthalene	107	Yes
2-Nonanone	180	Yes
Tetrahydrolinolool	202	Yes
Linalool	240	Yes
Dihydromyrcenol	307	Yes
Nonanal	502	Yes
Ethyl Undecylenate	624	Yes
Decanal	942	Yes
α -ionone	1,692	Yes
γ -methyl ionone	1,677	Yes
1-nonanol	24,164	Yes
Methyl Linolenate	771,304	Yes
Methyl Linolate	61149428	Yes

EXAMPLE 2

5 As noted in Table 1 of the specification, perfume may have various properties depending on hydrophobicity (oil/water partition coefficient), volatility (volatility constant), etc.

10 Figures 2(a) – 2(c) show calculated fragrance concentration profile in the headspace for three different types of perfume in a 5% surfactant product containing surfactant system with a CMC of 0.5% (i.e., sodium laurate). Hence, maximum fragrance burst is obtained with 10 dilutions (around CMC). Figure 2(a) is “type 2” perfume (high hydrophobicity, low volatility) where one would expect high burst at CMC; Figure 2(b) is “type 3” perfume (high hydrophobicity, high volatility), where burst at dilution is less than for Type 2; and Figure 2(c) is “type 4” perfume (low hydrophobicity; low volatility), where no “burst” would be expected.

15 In general, fragrances with high “perfume burst index” (PBI) will produce much larger burst upon dilution than those with medium PBI. Perfume molecules with PBI less than about 3 will essentially produce no perceivable enhancement in smell on dilution and simply show standard dilution curve with decreasing perfume concentration in headspace (see Figure 2(c)).

20 In general, it may be noted that one key to “burst” is that the perfume must have high hydrophobicity.

EXAMPLE 3

5

Several different fragrance molecules were tested experimentally to validate the theoretical predictions. Limonene, p-cymene, α -pinene, tetrahydrolinolool, α -ionone, γ -methyl ionone, benzyl acetate and 2-phenoxyethanol were tested in a 5% sodium laurate shower product and diluted a number of times through the CMC (Figure 3). The PBI of limonene is 12, p-cymene is 12.2, α -pinene is 20, tetrahydrolinolool is 202, α -ionone is 1692, γ -methyl ionone is 1677, benzyl acetate is -819 and 2-phenoxyethanol is -156. As predicted, limonene, p-cymene and α -pinene with a moderate positive PBI show moderate fragrance bursts of ~95%, 120% and 50% respectively. Tetrahydrolinolool, α -ionone and γ -methyl ionone with large PBI values give quite large fragrance bursts of 320%, 430% and 460%, while benzyl acetate and 2-phenoxyethanol with low, negative PBI values show no fragrance burst at all.

10
20
30
40
50
60
70
80
90
100
110
120
130
140
150
160
170
180
190
200
210
220
230
240
250
260
270
280
290
300
310
320
330
340
350
360
370
380
390
400
410
420
430
440
450
460
470
480
490
500

EXAMPLE 4

5 Besides simply obtaining a “fragrance burst” upon dilution of these surfactant
containing products, one can also produce a change in fragrance note. Many perfumes
are a combination of many different fragrance molecules. Depending on the
combination of molecules and the amount of each fragrance component in the mixture,
one can obtain both a “fragrance burst”, i.e. an overall increase in fragrance
10 concentration in the air around the product, and/or a change in fragrance note, i.e. a
change from a “floral/green” note to a citrus note. The reason this change in fragrance
note can be obtained is because upon product dilution, those molecules with a PBI
greater than about 3 will actually increase in concentration while those compounds with
a PBI less than about 3 will simply dilute and decrease in concentration. Therefore the
concentration of the different fragrance components smelled with product use will
change from that smelled over the fully formulated “original” product and hence the
overall character or “note” of the fragrance will be modified.

A few examples of fragrance mixtures that would be predicted to change note
upon product dilution or use are shown in Tables 4-5. Figure 4 shows examples of two
2-component fragrances which would definitely change note upon dilution as one
component increases in concentration and the other decreases.

Table 4: Perfume that Could Change "Note" on Dilution in Surfactant Containing Product

Fragrance Component	PBI	Dilution Behavior
Benzyl Acetate	-881	Decrease headspace (HS)
Cinnamyl Acetate	-43.3	Decrease HS
Limonene	12	Increase HS (50%)
Dihydromyrcenol	86	Increase HS (>3 times)
a-ionone	366	Increase HS (>3 times)

Table 5: Perfume that Could Change "Note" on Dilution in Surfactant Containing Product

Fragrance Component	PBI	Dilution Behavior
2-heptanone	-153	Decrease HS
Menthone	-71.4	Decrease HS
Isoeugenol	58	Increase HS (>3 times)
Tetrahydrolinalool	202	Increase HS (>3 times)
Linalool	240	Increase HS (>3 times)

EXAMPLE 5

5 It is also possible to take advantage of the technology of the invention to provide the introduction of a smell during dilution of the surfactant containing product. This can be achieved by formulating a fragrance ingredient into the product at a concentration just below the odor threshold. Therefore this component will not be perceived by the consumer in the original product, but as dilution occurs, if the PBI is large enough to supply a concentration of that component into the headspace above the product to exceed the odor threshold, the fragrance molecule will now be perceived upon use.

10 This concept can be used to provide a product that is essentially "fragrance-free", but produces a light fragrance with product dilution. This concept can also be utilized to change fragrance note, but providing a new smell upon product dilution to combine with the other perfume molecules to produce a new overall perfume mixture with use.

EXAMPLE 6

As suggested by the theoretical predictions, formulation factors also play a role in controlling the fragrance burst from the surfactant containing product in use. Two very important factors are the surfactant concentration in the original product and the CMC of the surfactant mixture used in the formulation.

The effect of surfactant concentration can be predicted from the theoretical calculations. The charts in Figure 5 show the result of two different perfumes with PBI of ~700 and ~12 with varying surfactant concentration (surfactant CMC = 0.005 wt/wt). As the graphs indicate, with a lower surfactant concentration in the original formulation, the initial fragrance concentration above the product will be higher (e.g., because fewer fragrance molecules are in surfactant micelles) and maximum fragrance burst can be reached with fewer dilutions. This is especially important, because the actual amount of dilution that typically occurs during product use is variable depending on the type of product and the consumer's habits. If the fragrance burst occurs with minimal dilution, the effect is more likely to be noticed by the product user. The other distinct advantage of products with low surfactant levels is that the absolute amount of fragrance available during the fragrance burst is greater, therefore the consumer will experience more fragrance during product use.

EXAMPLE 7

Experiments performed on actual perfumes in products with different surfactant concentrations are shown in Figure 6. Limonene (PBI=12.2) was tested in shower liquids containing sodium laurate at 5, 10, 15 and 20 wt.%, γ -Methyl Ionone and menthone were tested in sodium laurate at 5, 10 and 15 wt.% (Figure 6). As noted in the theoretical calculations in Example 6, the maximum fragrance burst is reached with fewer dilutions in the 5% soap product and the overall amount of fragrance available during the burst is also greater for the 5% product.

Thus, as seen, lower surfactant concentration is highly beneficial.

EXAMPLE 8

5 As noted in Examples 2 and 6, another variable that affects the fragrance burst
properties of a formulation is the critical micellization concentration (CMC) of the
surfactant system. The CMC is actually used to calculate the PBI for various fragrance
molecules. As the CMC of the surfactant system is decreased, the PBI values for
different fragrance molecules also decreases as shown in Table 6. Therefore in
10 products with surfactant systems with low CMC values, the expected extent of the
fragrance burst for different fragrances will be less.

Table 6: Fragrance Burst Potential in Surfactants with Different CMC's

Fragrance	PBI CMC=0.005	Burst	PBI CMC=0.0025	Burst	PBI CMC=0.001	Burst
Benzyl Acetate	-819	No	-1694	No	-4319	No
Hexanal	-109	No	-268	No	-599	No
Methyl Benzoate	-13	No	-32	No	-89	No
2-Octanone	-5	No	-178	No	-697	No
Isoeugenol	58	Yes	21	Yes	-90	No
Hexyl Acetate	86	Yes	-119	No	-188	No
Napthalene	107	Yes	78	Yes	-10	No
2-Nonanone	194	Yes	-94	No	-190	No
Linanol	240	Yes	147	Yes	-133	No
Limonene	12	Yes	9	Yes	1	No
α -Ionone	1692	Yes	1261	Yes	-30	No
α -Pinene	20	Yes	19	Yes	16	Yes
Tetrahydrolinolool	202	Yes	183	Yes	127	Yes
Dihydromyrcenol	307	Yes	267	Yes	145	Yes
Ethyl Undecylenate	624	Yes	557	Yes	357	Yes
Decanal	942	Yes	916	Yes	847	Yes
Methyl linolenate	771304	Yes	771283	Yes	771222	Yes

As expected, the lower the CMC of the surfactant systems, the lower the fragrance burst potential. Also, since the CMC is lower and the maximum burst is found around the CMC, as one moves to products with lower CMC's, the consumer will have to dilute the product more to actually experience higher fragrance concentrations.

EXAMPLE 9

Limonene and γ -Methyl Ionone were tested in two different shower liquid samples. One product was formulated with 5% sodium laurate (CMC = 0.005 wt/wt) and the second with 5% sodium laureth sulfate (CMC = 0.0011 wt/wt) (Figure 8). These results validate the theoretical results in Figure 7. The product with the lower CMC gives a lower overall fragrance burst and more dilutions are required to reach the maximum burst potential.

EXAMPLE 10

A typical fully formulated perfume used in a commercial product, e.g., personal wash formulation, usually is a multi-component composition in which the properties of each perfume component can vary dramatically. A typical perfume mixture in a 5 wt.% sodium laurate formulation was tested for its “perfume burst” behaviour upon dilution. The PBI values of the components in this perfume mixture vary from a large negative number (e.g., benzyl acetate, PBI = -819; PEA, PBI = -200 in sodium laurate solution) to a large positive number (e.g., gamma-methyl-ionone, PBI = 1,677 in sodium laurate solution). The dilution behaviour of each component in this perfume mixture is shown in Figure 9 (the graph represents a normalized GC measurement so the amount of burst of each molecule is clearly indicated). In this perfume mixture of twelve components, the dilution behaviour of each component is different. Each component has a unique dilution curve (GC headspace vs. dilution factor) governed by the factors (hydrophobicity, volatility, etc) discussed previously. For instance, for those perfumes with a negative PBI (e.g., benzyl acetate), the perfume headspaces decrease upon dilution. For those perfumes with a moderate positive PBI (e.g., limonene, PBI = 12; p-cymene = 12.2 in sodium laurate solution), the perfume headspaces increase 1-2 times upon dilution. For those fragrance components with a very large positive PBI (e.g., gamma-methyl-ionone), the “perfume burst” can be as high as seven times. The results shown in Figure 9 indicated that in a fully formulated perfume, in terms of dilution behaviour, the performance of each component is mostly determined by its own individual physical properties.

EXAMPLE 11

Trained sensory panel analyses were carried out to validate the “perfume burst” phenomena. A group of 20 to 30 expert sensory panellists were asked to rate the intensity of the fragrance over formulations and anchor their scores to a perfume standard supplied for each test. Formulations containing a single perfume (gamma-methyl-ionone) and multi-component perfume were studied in two separate panels. The multi-component perfume contains four perfumes of equal weight percentage, menthone, tetra-hydro-linalool, α -ionone, γ -methyl-ionone, that will all “burst” in a sodium laurate product, but will not “burst” (e.g., menthone, α -ionone) or will give a moderate “burst” (e.g., tetra-hydro-linalool, γ -methyl-ionone) in an SLES product with dilution. Each sample was tested in duplicate and the sample information and the panel study results are shown in Tables 7 and 8, and Figures 10 and 11. As shown in Tables 7 and 8, perfumes at two different initial concentrations were tested in the sodium laurate solution. The scores shown in Figures 10 and 11 are the panel averages for each sample.

The results of the panel analyses indicate that as expected, both the single perfume and the perfume mixture (Figure 10 and 11) in a sodium laurate product (a surfactant of high CMC, CMC = 0.5 wt.%), have higher sensory scores for the 10 times diluted solution compared to the original undiluted formulations. In the SLES formulations (a surfactant of relatively low CMC, CMC = 0.1 wt.%), both single perfume and perfume mixture (Figure 10 and 11) have lower sensory scores for the 10 times diluted solution compared to the original undiluted product. These results are consistent with the discussion and data presented in Example 8, which indicate that a surfactant of low CMC lowers the PBI of the perfume and can greatly reduce the expected “perfume burst” upon dilution.

The panel studies indicate that this “perfume burst” upon dilution can be perceived quite easily by the human nose if the formulation is designed according to the rules listed in this invention.

5 Table 7: Sample Information of the Single Perfume (γ -methyl-ionone) System for Panel Study

Sample Name	Information
1	0.0649 wt.% γ -methyl-ionone in 4.5 wt.% sodium laurate
2	10 times dilution of 0.0649 wt.% γ -methyl-ionone in 4.5 wt.% sodium laurate
3	0.2193 wt.% γ -methyl-ionone in 4.5 wt.% sodium laurate
4	10 times dilution of 0.2193 wt.% γ -methyl-ionone in 4.5 wt.% sodium laurate
5	0.2200 wt.% γ -methyl-ionone in 4.5 wt.% SLES
6	10 times dilution of 0.2200 wt.% γ -methyl-ionone in 4.5 wt.% SLES

Table 8: Sample Information of the Multi-Component Perfume (menthone, tetra-hydro-linalool, α -ionone, γ -methyl-ionone) System for Panel Study

Sample Name	Information
7	0.2404 wt.% perfume mixture in 5 wt.% sodium laurate
8	10 times dilution of 0.2404 wt.% perfume mixture in 5 wt.% sodium laurate
9	0.6022 wt.% perfume mixture in 5 wt.% sodium laurate
10	10 times dilution of 0.6022 wt.% perfume mixture in 5 wt.% sodium laurate
11	0.2433 wt.% perfume mixture in 5 wt.% SLES
12	10 times dilution of 0.2433 wt.% perfume mixture in 5 wt.% SLES

EXAMPLE 12

There are any number of surfactant containing formulations that should provide a "perfume burst" in use. These include personal wash products (i.e. shower gels, soap bars), shampoos, household cleaners, light duty detergents, fabric washing products, etc. All of these products undergo some dilution with use which could potentially cause release of fragrance molecules from the surfactant micelles and provide a "burst" of fragrance intensity. As explained in the previous examples, the CMC of the surfactant mixture, the amount of surfactant in the product and the perfume ingredients used in the fragrance formulation all contribute to the type and intensity of the "perfume burst" during product use. The following formulations are just a few examples of products that would be expected to provide a "fragrance burst" or change in fragrance note, assuming the perfume was properly designed (i.e. a fragrance containing a fraction of components with high PBI values).

Table 9

Formulation Ingredients (%)	Form. #1	Form. #2	Form. #3	Form. #4	Form. #5	Form. #6	Form. #7	Form. #8
Sodium Laurate	10	5	8	4			4	5
TEA Laurate		5			5			
Sodium Laureth Sulfonate			2		4			8
Sodium Lauryl Sulfate						8		
Alkylpoly Glucoside				7				
Nonionic Polymeric Surfactant Containing Mixture of Alkylene Oxides		2		5				
Caprylamido Propylbetaine	1				1	2		1
Sodium Cocoylisethionate							10	
Lauric Acid	3		2	2			3	
Amine Oxide						1		
Xanthan Gum		1						0.8
Jaguar S13 (Cationic Polymer)					0.4			
Carbopol Polyacrylate (e.g., Structuring Agent)	0.6		0.6					
Polymer JR (Cationic Polymer)							0.3	
Triclosan		0.5						
Trichlorocarbanilide				0.5				
Propylene Glycol								10
Glycerine							30	
Vitamin E					0.05			
Perfume	1	1	1.3	1	1	1.2	0.8	1
Water	To 100	To 100	To 100	To 100	To 100	To 100	To 100	To 100

As noted, these are examples of the many formulations in which the perfume
 5 burst concept can be used.

EXAMPLE 13

The deposition of the fragrance components onto a surface (e.g., skin) upon dilution of a surfactant containing product is also closely related to the thermodynamic characteristics of the perfume (e.g., oil/water partition coefficient, ϕ , and volatility constant, K) and the formulation (e.g., the surfactant CMC). For a perfume-containing composition that has a high PBI, the perfume headspace will increase upon dilution due to the increase of the perfume concentration in the continuous phase caused by surfactant demicellization. This increase in perfume concentration in the continuous phase will also favor the deposition of the perfume onto the surface that the product comes into contact with due to the enhanced chemical potential of the perfume in the continuous phase which is in direct contact with the surface. Since perfume deposition is directly correlated with perfume burst, the PBI can also be used to predict fragrance delivery to a substrate during product use. Therefore, for a certain perfume, the surfactant systems which facilitate perfume burst upon dilution can also enhance the deposition of perfume onto the surface in contact, compared to the surfactant systems which less favor the burst.

The deposition of a Type 2 perfume which typically have high PBI values was tested in two surfactant systems with different CMCs (sodium laurate, CMC=0.5 wt. %, and SLES, CMC=0.1 wt. %) on a synthetic skin substrate (e.g., Vitro-Skin®). The recovery of the perfume, which is the ratio of the amount of perfume extracted after deposition to that originally dosed onto the surface, is used to reflect the deposition efficiency of the perfume. The results of the recovery of γ -methyl-ionone are plotted vs. the dilution factor for both sodium laurate and SLES surfactant systems (Figure 12). The recovery of γ -methyl-ionone from the skin washed with the undiluted sodium laurate and SLES surfactant systems (the surfactant concentration in both systems is 5%) is not significantly different. Upon dilution, the recovery of the perfume is enhanced in both surfactant systems, however, the sodium laurate surfactant system, which has a higher PBI due to the higher CMC of the surfactant, provides a much higher deposition efficiency upon dilution than the SLES surfactant system. These results correlate as expected with the perfume burst results.

EXAMPLE 14

5 The deposition of a multi-component "burst" perfume composed of perfumes with relatively high PBI's (i.e. high oil/water partition coefficient (ϕ) and low volatility constant (K)) was tested on vitro-skin from sodium laurate surfactant system (high CMC) and SLES surfactant system (low CMC). The results were plotted in Figure 13. It was found that the recoveries of the perfumes from this mixture after deposition from the undiluted sodium laurate and SLES surfactant systems (the surfactant concentration in both systems is 5%) are very similar. An enhanced recovery of the perfume in both diluted surfactant systems (sodium laurate and SLES) is observed compared with that of the perfume in the undiluted surfactant systems. However, after 10 times dilution, the perfume recoveries were higher in sodium laurate surfactant system than that in SLES. These results again correlate with the perfume burst measurements, indicating that the PBI can be a useful tool for predicting fragrance deposition from surfactant containing products.